

The Synergistic (MARATHON) Effect of Combined Methamphetamine with Sexual Stimulant Drugs on Increasing the Likelihood of High-Risk Sexual Behaviors

Seyed Mehdi Hosseinifard PhD¹, Alireza Ahmadian MSc², Neda Smaelifar MSc³

Original Article

Abstract

Background: Chronic drug abuse and sexual dysfunction specifically erectile dysfunction may lead drug abusers to seek over-the-counter or non-prescription medications, out of which Sildenafil citrate, sold as the trade name of Viagra® can be considered as a prime and important treatment. Therefore, the research purpose was to draw a comparison and review the role of methamphetamine abuse and sildenafil use in increasing the likelihood of high-risk sexual behaviors (both concomitant and non-concomitant use).

Methods: Hence, a total of 40 patients diagnosed with methamphetamine abuse were recruited through the administration of structured clinical interview for DSM-IV (Diagnostic and Statistical Manual of Mental Disorders, 4th Edition), through purposive sampling and subsequent to being qualified in accordance with the selection criteria by psychologists and general practitioners. All the 40 drug abusers (20 methamphetamine abusers with concomitant use of Aphrodisiac drugs (sexual stimulant pills) and 20 methamphetamine abusers) described their sexual risk behaviors subsequent to the drug use.

Findings: Supported the between-group difference that is to say that, the group with concomitant methamphetamine abuse differed significantly in all the items when compared with the control group. However, this group scored lower on the item of sexual intercourse with drug addicted prostitutes using condom and both groups demonstrated high pick on this item.

Conclusion: Overall, the concomitant methamphetamine chronic abuse with sexual stimulant drugs generates Aphrodisiac drugs impulses and is found to be related to higher frequencies of sexual risk behaviors and sexual intercourse with addicted prostitutes.

Keywords: Methamphetamine, Aphrodisiac drugs, High-risk sexual behaviors

Citation: Hosseinifard SM, Ahmadian A, Smaelifar N. **The Synergistic (MARATHON) Effect of Combined Methamphetamine with Sexual Stimulant Drugs on Increasing the Likelihood of High-Risk Sexual Behaviors.** *Addict Health* 2014; 6(3-4): 112-8.

Received: 03.04.2014

Accepted: 16.06.2014

1- Assistant Professor, Department of Psychiatry, School of Medicine, Kerman University of Medical Sciences, Kerman, Iran

2- PhD Student, Department of Family Counseling, School of Humanities, Kharazmi University, Tehran, Iran

3- PhD Student, Department of Family Counseling, School of Humanities, Kharazmi University AND Young Researchers and Elite Club, Islamic Azad University, Roudheh Branch, Tehran, Iran

Correspondence to: Seyed Mehdi Hosseinifard PhD, Email: smhf56@gmail.com

Introduction

Methamphetamine (methylanphetamine) can be classified into the drug class of central nervous system stimulants and is a psychostimulant and psychoactive drug with numerous slang terms that vary from region to region such as ice, clouds crystal or glass.^{1,2} Methamphetamine triggers a cascading release of dopamine in the brain mechanism, and its psychological effects can be marked by dramatic mood swings from euphoria and elation to irritability.^{3,4}

Methamphetamine is most structurally similar to amphetamine and most commonly, it is found as a colorless, bitter and non-smelling crystalline solid. Long-term use can have irritability, psychomotor agitation, increased risk of Parkinson's disease, chronic depression and sexual impulsiveness.^{5,6} In addition, delusions of grandiosity, hallucinations, hyperactivity, excessive feelings of power and invincibility, repetitive and obsessive behaviors, paranoia are some of the physical and psychological symptoms of methamphetamine abuse with chronic use and/or high dose.⁷ At the start of amphetamine abuse, the abuser may exhibit exaggerated and changing sexual attitude and feelings, unprotected and unsafe sex acts, strong and uncontrollable sexual impulses, frequent change of sexual partners and irresponsible sexual behavior.⁸ The abuser is not able to perceive wrong-right behaviors while having sex and may display impulsive or reckless behavior. Over time, substance abuse and dysfunction in sexual performance specifically erectile dysfunction may give rise to the use of illegal aphrodisiac drugs (sexual stimulant pills). Overall, methamphetamine use is found to be related to higher frequencies of high-risk sexual activities.^{9,10}

Sildenafil citrate, sold as Viagra®, revatio and under various other trade names, is a drug used to treat erectile dysfunction and pulmonary arterial hypertension. It acts by inhibiting cyclic guanosine monophosphate (cGMP)-specific phosphodiesterase Type 5, an enzyme that promotes degradation of cGMP, which regulates blood flow in the penis. Since becoming available in 1998, sildenafil has been the prime treatment for erectile dysfunction.¹¹ Patterson and Semple,¹² Crosby and Mettey,¹³ Halkitis and Green,¹⁴ and Spindler et al.² found that illegal and even legal

stimulant drugs in substance abusers, allows for further probability of sexual risk behaviors. This negative combined synergistic effect is called sex-drug cocktail in the west. Methamphetamines entail higher risk of shaping and reinforcing sexual risk behaviors when compared to other substances such as hashish, opium, crack, heroin, Ritalin and cocaine. This seems to be due to the neurochemical synergistic reaction. One out of five HIV (human immunodeficiency virus)-positive participants was methamphetamine abuser in a longitudinal study done in Los Angeles.¹⁵

The research findings indicate that methamphetamine abuse heighten the risk of engagement in unprotected intercourse and riskier sexual behaviors.¹⁶ The present research is an attempt to shed light on the role of methamphetamine abuse and sildenafil (Viagra®) use in increasing the likelihood of high-risk sexual behaviors (concomitant and non-concomitant use).

Methods

A total of 40 patients diagnosed with methamphetamine abuse were recruited through the administration of structured clinical interview for DSM-IV (SCID-I). The diagnosed abusers were selected through purposive sampling.^{17,18} Methamphetamine abusers with concomitant and non-concomitant use of aphrodisiac drugs (sexual stimulant pills) were selected from among the patients who were referred to counseling and harm reduction centers (triangular counseling clinics of AIDS (acquired immune deficiency syndrome), substance abuse and sexually transmitted diseases treatment) in Tehran, Iran. Subsequent to being qualified in accordance with selection criteria by psychologists and general practitioners, all the 40 drug abusers (20 methamphetamine abusers with concomitant use of aphrodisiac drugs (sexual stimulant pills) and 20 methamphetamine abusers) described their sexual risk behaviors subsequent to the drug use.

The inclusion criteria included chronic methamphetamine abuse (over 3.5 years) at the present time, lack of multiple substance abuse during the previous addiction or earlier (the main previous addiction had started with methamphetamine), aged between 25 and 40 years and the minimum level of secondary education to fill out the required forms and checklists. The exclusion criteria were having an

acute physical illness and concurrent use of other specific substances before or during the study.

To achieve this goal, The SCID-I Axis 1 disorders, was conducted by a clinical psychologist. Subsequent to the initial diagnosis of chronic methamphetamine abuse, the above criteria were considered to select the eligible representative sample (Table 1). Comorbidity with other disorders is usually observed among the patients with substance abuse.¹⁹ Ladouceur et al.²⁰ have noticed the same problem in their clinical studies. According to this approach and in order to overcome this problem we have selected individuals with the primary diagnosis of chronic methamphetamine addiction and concomitant sexual abuse drugs. Therefore, 40 controlled patients were enrolled subsequent to the selection of the eligible sample.

SCID-Clinical version (CV): The SCID-I Axis 1 disorders: SCID-CV, which was developed by Spitzer²¹ was specially designed as an adaptation of the SCID that is intended to introduce the benefits of structured interviewing into clinical and research setting. SCID-CV helps not to resort to the lengthier and more complex process used principally in research studies and dispose of other diagnostic categories which are not related to the study.²² The SCID-CV is divided into six self-contained modules covering: module A: mood episode, module B: psychotic symptoms, module C: psychotic disorders, module D: mood disorders, module E: substance use disorders, module F: anxiety and other disorder, also included in module F are disorders without diagnostic criteria such as agoraphobia, social phobia, specific phobia, etc. The SCID-CV may be administered to either psychiatric or general medical patients. It is most appropriate for adults (18 years and over), but with slight modification, may be used with adolescents.²³

This instrument can't be carried out with patients who suffer from serious cognitive disorders or exhibit severe and distressing symptoms of psychotic disorders and having

acquired at least 8 years of education is required for full understanding.²⁴ Tran and Haaga²⁵ have reported the results indicate moderate to good reliability among formulations constructed by teams of independent clinician that was equal to 0.60 based on a weighted kappa. A Persian version of this interview was administered on 229 individuals by Sharifi et al.²⁶ and the result of the kappa coefficient was over 0.60. In order to assess the validity of the instrument, Bakhtiyari²² asked a panel of judges formally to rate the validity, which was rated as having satisfactory and acceptable validity to be used in clinical setting. The test was also shown to have a high 1-week test-retest reliability ($r = 0.95$).

Substance Abusers' sexual behaviors questionnaire: this 12-item questionnaire was designed and developed by Vaziri and Lotfi Kashani²⁷ and measures sexual desire and performance, peak sexual demands and sexual risk behaviors. Each dimension is covered with 4 questions. This questionnaire was administered on 30 substance abusers and its test-retest reliability with 15 days of the time interval was equal to 0.76. Each question has been designed in a way to provide a total score for each dimension, and it is possible to compare individuals in each of the possible responses. This questionnaire is based on a four-point scale ranging from 0 "never" to 3 "always" and can be marked verbally or in written form.²⁴ With regard to the research condition and patients' chronic problem and subsequent to the required evaluations, the short 5-item questionnaire was designed and completed by the respondents (Table 1).

Results

As seen, the sample population was selected based on the demographic characteristics and comparable clinical severity in terms of age, sex and duration of abuse. The mean age and the mean of the total score of the first group with methamphetamine abuse and taking sexual stimulant drug were 34.50 and 2.24, respectively.

Table 1. The aim of questionnaires' administration and their timing

| Questionnaire | Aim | Timing |
|----------------|--|---|
| SCID-I | Clinical differential diagnosis of methamphetamine abuse | Prior to patient selection and during the screening phase |
| Oral interview | Self-report diagnosis of methamphetamine abuse | outlining the baseline level |

SCID-I: Structured clinical interview for DSM-IV (Diagnostic and Statistical Manual of Mental Disorders, 4th Edition)

Moreover, the mean age and the mean of the total score of the second group with methamphetamine abuse were 35.00 and 0.97, respectively (Table 2).

Table 3 illustrates the means of the items in the questionnaire in two groups. As observed, the first group with concomitant abuse of methamphetamine and sexual stimulant drug scored higher on the items of sexual power and performance with substance abuse, rapid, sharp and frequent increase of sexual behavior, sexual intercourse with strangers without condom, anal and oral sex and masturbation and sexual intercourse with drug addicted prostitutes with condom.

Moreover, the group with concomitant abuse of methamphetamine and sexual stimulant drug scored higher on the items of sexual power and performance with substance abuse, rapid, sharp and frequent increase of sexual behavior, sexual intercourse with strangers without condom, anal and oral sex and masturbation and sexual intercourse with drug addicted prostitutes with condom (Table 3).

Discussion

The data presented in figure two demonstrate the between-group difference on all the items specifically the fourth one. Aphrodisiac drugs (sexual stimulant pills) users showed a sharp increase of sexual stimulations on all the items when compared with the non-using group. Overall, as observed, both groups showed tendency to have sexual intercourse with drug addicted prostitutes using condoms.

The sexual desire in single methamphetamine

drug abuse was only slightly less than concomitant methamphetamine abuse and Aphrodisiac drugs (sexual stimulant pills). However, high-risk sexual behaviors are due to their decreasing and unacceptable level of socio-sexual networks, financial poverty, and emotional strains. Unpleasant and visible symptoms of methamphetamine addicted individuals result in the elimination of their names from the customers' list of healthier and more high-class prostitutes. Therefore, they seek for street and low-cost prostitutes who are typically addicted. Concomitant methamphetamine abuse with sexual stimulant drugs generates sexual impulses that reinforce high-risk sexual behaviors with a network of sexual partners who are likely of contracting a sexually transmitted infection (STI) even with using a condom. Providing and distributing condoms can help these individuals recognize the benefits of the less riskier sexual activities and reduced likelihood of contracting HIV and STI; however, some of them cannot control themselves in sexual intercourse with partners who are highly likely of being HIV- or STI-positive and their preference was obvious on the 5th item of the questionnaire which indicated their tendency in using condom in high-risk sexual activities (Figure 1).

Conclusion

Overall, it can be concluded that the concomitant methamphetamine chronic abuse with Aphrodisiac drugs (sexual stimulant pills) generates aphrodisiac drugs impulses and is found to be related to higher frequencies of sexual risk behaviors and sexual intercourse with addicted

Table 2. Patients' demographic characteristics

| Age range (year) | Sex | No. | Duration | Sample | Mean of total score |
|------------------|------|-----|------------------------|--|---------------------|
| 34.5 | Male | 20 | 4 years, approximately | Methamphetamine abuse and sexual stimulant drugs | 2.24 |
| 35.0 | Male | 20 | 4 years, approximately | Methamphetamine abuse | 0.97 |

Table 3. Comparisons of groups' mean scores on the five items of substance abusers' sexual behaviors questionnaire

| Item | Mean | |
|---|-------------------|-----------------------|
| | Concomitant abuse | Non-concomitant abuse |
| Increase of sexual power and performance with substance abuse | 2.30 | 0.73 |
| Rapid, sharp and frequent increase of sexual behavior | 2.40 | 0.83 |
| Sexual intercourse with strangers without condom | 1.90 | 0.90 |
| Anal and oral sex and masturbation | 1.80 | 0.70 |
| Sexual intercourse with drug addicted prostitutes with condom | 2.80 | 1.70 |




Figure 1. Comparison of high-risk sexual behaviors in two groups of non-concomitant methamphetamine abuse and concomitant methamphetamine abuse and sexual stimulant drugs

prostitutes. These findings point to sexual marathons as a possible contributor to rising HIV rates in the combined methamphetamine drug and sexual stimulant drugs.

Conflict of Interests

The Authors have no conflict of interest.

Acknowledgements

It would be the authors' pleasure to express their gratitude to Dr. Afsoon Rahimian Shad, the Master of Public Health (MPH).

References

1. Swearingen SG, Klausner JD. Sildenafil use, sexual risk behavior, and risk for sexually transmitted diseases, including HIV infection. *Am J Med* 2005; 118(6): 571-7.
2. Spindler HH, Scheer S, Chen SY, Klausner JD, Katz MH, Valleroy LA, et al. Viagra, methamphetamine, and HIV risk: results from a probability sample of MSM, San Francisco. *Sex Transm Dis* 2007; 34(8): 586-91.
3. American Psychiatric Association, Task Force on Nomenclature and Statistics. *Diagnostic and Statistical Manual of Mental Disorders*. Arlington, TX: American Psychiatric Association; 1994.
4. Sadock BJ, Kaplan HI, Sadock VA. Kaplan & Sadock's Synopsis of Psychiatry: Behavioral Sciences/clinical Psychiatry. Philadelphia, PA: Lippincott Williams & Wilkins; 2007.
5. Reback CJ. The social construction of a gay drug: Methamphetamine use among gay and bisexual males in Los Angeles [Online]. [cited 1997]; Available from: URL: http://www.uclaisap.org/assets/documents/final-report_cjr_1-15-04.pdf
6. Semple SJ, Patterson TL, Grant I. Motivations associated with methamphetamine use among HIV⁺ men who have sex with men. *J Subst Abuse Treat* 2002; 22(3): 149-56.
7. Schultz W. Potential vulnerabilities of neuronal reward, risk, and decision mechanisms to addictive drugs. *Neuron* 2011; 69(4): 603-17.
8. Dalley JW, Everitt BJ, Robbins TW. Impulsivity, compulsivity, and top-down cognitive control. *Neuron* 2011; 69(4): 680-94.
9. Belenko S, Fabrikant N, Wolff N. The Long Road to Treatment Models of Screening and Admission into Drug Courts. *Criminal Justice and Behavior* 2011; 38(12): 1222-43.
10. Ahmadian AR, Shokri B, Smaeeli N. Pictorial encyclopedia of substance abuse. Tehran, Iran: Sayeh Nima Publication; 2014. p. 114-63.

11. Mansergh G, Shouse RL, Marks G, Guzman R, Rader M, Buchbinder S, et al. Methamphetamine and sildenafil (Viagra) use are linked to unprotected receptive and insertive anal sex, respectively, in a sample of men who have sex with men. *Sex Transm Infect* 2006; 82(2): 131-4.
12. Patterson TL, Semple SJ. Sexual risk reduction among HIV-positive drug-using men who have sex with men. *J Urban Health* 2003; 80(4 Suppl 3): iii77-iii87.
13. Crosby R, Mettrey A. A descriptive analysis of HIV risk behavior among men having sex with men attending a large sex resort. *J Acquir Immune Defic Syndr* 2004; 37(4): 1496-9.
14. Halkitis PN, Green KA. Sildenafil (Viagra) and club drug use in gay and bisexual men: the role of drug combinations and context. *Am J Mens Health* 2007; 1(2): 139-47.
15. Rosenblatt S. More gay men using meth, study finds. Use of the drug, which is associated with HIV transmission, has surged since 2005, according to data collected by a nonprofit agency. *Los Angeles Times* 2007 Apr 11.
16. Fisher DG, Reynolds GL, Jaffe A, Johnson ME. Reliability, sensitivity and specificity of self-report of HIV test results. *AIDS Care* 2007; 19(5): 692-6.
17. Kazdin AE. Mediators and mechanisms of change in psychotherapy research. *Annu Rev Clin Psychol* 2007; 3: 1-27.
18. Gall MD, Gall GP, Borg WR. *Educational Research: An Introduction*. Boston, MA: Pearson/Allyn & Bacon; 2007.
19. Ball SA, Young JE. Dual Focus Schema Therapy for personality disorders and substance dependence: Case study results. *Cognitive and Behavioral Practice* 2000; 7(3): 270-81.
20. Ladouceur R, Gosselin P, Dugas MJ. Experimental manipulation of intolerance of uncertainty: a study of a theoretical model of worry. *Behav Res Ther* 2000; 38(9): 933-41.
21. Spitzer RL. *User's Guide for the Structured Clinical Interview for DSM-III-R: SCID*. Arlington, TX: American Psychiatric Press; 1990.
22. Bakhtiyari M. *Mental disorders in patients with body dysmorphic disorder*. [Thesis]. Tehran, Iran: Tehran Psychiatric Institute; 2000. [In Persian].
23. Barlow DH. *Clinical Handbook of Psychological Disorders, Fifth Edition: A Step-By-Step Treatment Manual*. New York, NY: Guilford Publications; 2014.
24. Bakhshipour A, Mohammadkhani S. *Practical Guide for Prevention and Treatment of Substance Abuse*. Tehran, Iran: Ministry of Health and Medical Education. Office of Substance Abuse Prevention and Treatment; 2005. [In Persian].
25. Tran GQ, Haaga DAF. Coping Responses and Alcohol Outcome Expectancies in Alcohol Abusing and Nonabusing Social Phobics. *Cognitive Therapy and Research* 2002; 26(1): 1-17.
26. Sharifi V, Asadi SM, Mohammadi MR, Amini H, Kaviani H, Semnani U, et al. Reliability of the Persian version of the structured clinical diagnostic interview based on DSM-IV for SCID. *Advances in Cognitive Science* 2004; 6(1-2): 10-22. [In Persian].
27. Vaziri S, Lotfi Kashani F. Effects of Methamphetamine and Narcotics on the Increase of Libido and Reckless Sexual Behavior. *Thought and Behavior in Clinical Psychology* 2010; 4(15): 81-91. [In Persian].

اثر ترکیب همافزا (ماراتن) سوء مصرف همزمان مت‌آمفتامین با داروهای محرک جنسی در افزایش بروز رفتارهای پرخطر آمیزشی

دکتر سید مهدی حسینی فرد^۱، علیرضا احمدیان^۲، ندا اسماعیلی فرد^۳

مقاله پژوهشی

چکیده

مقدمه: با مرور زمان و سیر مزمن سوء مصرف مواد و ناتوانی در برخی حوزه‌های کارکرد جنسی به خصوص ناتوانی نعوظ، فرد سوء مصرف کننده اقدام به درمان‌های خودسرانه مصرف داروهای محرک جنسی می‌کند که یکی از مهم‌ترین این داروها سیلدنافیل سیترا (ویاگرا) نام دارد. مطالعه حاضر گزارش مختصری از بررسی نقش سوء مصرف مت‌آمفتامین و سیلدنافیل (ویاگرا) به تنهایی و در موقعیت ترکیب با هم در احتمال بروز رفتارهای جنسی پرخطر بود.

روش‌ها: ابزار SCID-I (Structured clinical interview for diagnostic and statistical manual of mental disorders) جهت تشخیص سوء مصرف مزمن مت‌آمفتامین استفاده گردید. معیارهای ورود به پژوهش و خروج از آن توسط روان‌شناس و پزشک مرکز انجام و ۴۰ آزمودنی سوء مصرف کننده مزمن وارد مرحله ثبت خط پایه شدند. هر ۴۰ آزمودنی (۲۰ نفر سوء مصرف توأم مت‌آمفتامین با سیلدنافیل و ۲۰ نفر هم سوء مصرف تنها با مت‌آمفتامین) به توصیف رفتارهای پرخطر جنسی خود پس از مصرف مواد پرداختند.

یافته‌ها: گروه سوء مصرف توأم با مت‌آمفتامین در همه بخش‌ها نسبت به گروه شاهد تفاوت معنی‌داری داشت که البته این تفاوت در بخش روابط با زنان روسپی معتاد با کاندوم کمتر بود و هر دو گروه در این گویه برافراشتگی زیادی داشتند.

نتیجه‌گیری: سوء مصرف توأم مت‌آمفتامین مزمن با داروهای محرک جنسی سبب تولید تکانه‌های تحریک جنسی می‌شود که علاوه بر تولید رفتارهای پرخطر نوعی ارتباط جنسی با شبکه‌ای از زنان مشکوک به آلودگی را (هرچند با استفاده از کاندوم) رقم می‌زند.

واژگان کلیدی: مت‌آمفتامین، داروی محرک جنسی، رفتار پرخطر جنسی

ارجاع: حسینی فرد سید مهدی، احمدیان علیرضا، اسماعیلی فرد ندا. اثر ترکیب همافزا (ماراتن) سوء مصرف همزمان مت‌آمفتامین با داروهای محرک جنسی در افزایش بروز رفتارهای پرخطر آمیزشی. مجله اعتیاد و سلامت ۱۳۹۳؛ ۶ (۳-۴): ۱۱۸-۱۱۲.

تاریخ پذیرش: ۹۳/۳/۲۶

تاریخ دریافت: ۹۳/۱/۱۴

۱- استادیار، گروه روانپزشکی، دانشکده پزشکی، دانشگاه علوم پزشکی کرمان، کرمان، ایران

۲- دانشجوی دکتری، گروه مشاوره خانواده، دانشکده علوم انسانی، دانشگاه خوارزمی، تهران، ایران

۳- دانشجوی دکتری، گروه مشاوره خانواده، دانشکده علوم انسانی، دانشگاه خوارزمی و باشگاه پژوهشگران و نخبگان، دانشگاه آزاد اسلامی، واحد رودهن، تهران، ایران

Email: smhf56@gmail.com

نویسنده مسئول: دکتر سید مهدی حسینی فرد